

# Comparing ICESat-2, SWOT, and GNSS-Reflectometry Water Surface Profiles over the Tonle Sap Lake

**Margaret Scott <sup>1</sup>, Alexa Angelina Putnam <sup>1</sup>, Rashmi Shah <sup>2</sup>,  
Cedric David <sup>2</sup>, Carolyn Roesler <sup>1</sup>, J. Toby Minnear <sup>1</sup>, and Y.  
Jade Morton <sup>1</sup>**

*<sup>1</sup>University of Colorado, Boulder*

*<sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena,  
CA, US*



margaret.scott-1@colorado.edu

# Motivation & Focus Area

- **Motivation:**

The impact of changing conditions (e.g., drought, damming) can be studied by looking at changes in the surface profile of inland water bodies measured by GNSS-Reflectometry


- **Focus Area: Tonle Sap Lake, Cambodia**

- Dry season (Nov-May):

- *Lake drains into the Mekong River*
- *Water Level ~2m; 2500 km<sup>2</sup> surface area*


- Rainy season (June-Oct):

- *Mekong floods-> Tonle Sap river reverses & fills the lake*
- *Water level rises to 10 m; 15000 km<sup>2</sup> surface area*



# Dataset 1: Spire grzRfl GNSS-R

- **Polar, ~500km Orbit**
- **Dual frequency GPS L1(1575.42 MHz) & L2 (1227.6 MHz)**
- **Grazing angle:**
  - 5-30deg elv. angle
- **Land collections Feb, 2021 – Present (our study is 2021-2023)**
- **50 Hz, L1B Product:**
  - Georeferenced excess carrier phase & power data
- **Carrier-phase height is relative**
  - Reference surface from DAHITI+EGM08



$$H = -\frac{\Delta\rho_{meas} - \Delta\rho_{mod} - B + \epsilon}{2 \sin(\theta)}$$



$$\Delta\rho_{meas} = \lambda_L(\phi_L^R - \phi_L^D) + \lambda_L n_L \quad \epsilon = \text{error}$$

$\Delta\rho_{mod}$  = modelled bistatic delay (geometric range, troposphere, ionospheric advance, carrier ambiguity)

C. Roesler, et al.,  
2022

# Dataset 2: ICESat-2

- **532 nm photon-counting with 6 laser beams:**
  - 3 strong, 3 weak
- **10 kHz PRF**
- **Absolute height measurement measuring 2-way time of flight**
- **ATL03 Product: georeferenced photon level data product**
- **Height measured as the median value between GNSS-R specular points (relative to EGM08)**



# Dataset 3: Surface Water and Ocean Topography (SWOT)

- **Launched 2022**
- **Primary Instrument KaRIN:**
  - Near-Nadir Ka-band Interferometric SAR
  - 35.75 GHz (Ka-band)
- **Data products used:**
  - PIXC
  - L2 rasterized product






Image Credit: E. Peral  
(2024)

# Why GNSS-R?

- **High spatial & temporal resolution**
- **Diverse look directions across lake**
- **Penetrates cloud and vegetation cover**




Accumulated Coherent Spire GNSS-R Tracks, 2021-2024






## Results: 1 Case Study

November 21, 2023 (ICESat-2), December 2, 2023 (GNSSR), & December 4, 2023 (SWOT)







November 21, 2023 (ICESat-2), December 2, 2023 (GNSSR), & December 4, 2023 (SWOT)



Date	Sensor	Open Water $\sigma_{rms}$ (cm)	Water under Vegetation $\sigma_{rms}$ (cm)	Open water R (GNSS,SWOT) ( $\alpha = 0.05$ )	Open water R (ICESat-2,SWOT) ( $\alpha = 0.05$ )	Open water R (GNSS,ICESat-2) ( $\alpha = 0.05$ )
November/ December, 2023	ICESat-2	1.6 cm	6.9 cm	0.91	0.94	0.93
	GNSS-R	2.6 cm	2.2 cm			
	SWOT	2.3 cm	4.6 cm			




## Interpreting Slope

Measurements important to freshwater resources are height (h), change of height of space (dh/dx), and change of height over time (dh/dt)




**HOWEVER!** It is well-established that geoid anomalies not captured at the resolution of geoid models such as EGM08 can exist over large lakes

# PIXC → L2 Rasterized data from SWOT 0.02° x 0.02° gridding

ICESat-2 (November 21, 2023), GNSS-R (December 2, 2023) and SWOT (December 4, 2024) Comparison



# Comparing across Distance, Height, and Time



- ICESat-2
- ★ GNSS-R
- SWOT





Motivation

Methods

Results

Summary

# Stable structure strongly suggestive of a geoid anomaly



- | Seasonally Inundated |                               |
|----------------------|-------------------------------|
| ●                    | SWOT 02-Aug-2023 09:16:57     |
| ●                    | SWOT 11-Sep-2023 16:56:55     |
| ●                    | SWOT 05-Oct-2023 09:06:55     |
| ●                    | SWOT 24-Oct-2023 17:40:15     |
| ●                    | SWOT 14-Nov-2023 05:09:08     |
| ●                    | SWOT 04-Dec-2023 16:25:31     |
| ●                    | SWOT 05-Feb-2024 18:40:24     |
| ●                    | SWOT 08-Apr-2024 20:11:06     |
| ●                    | SWOT 20-May-2024 21:37:47     |
| ★                    | GNSSR 12-02-2023 14:35:22     |
| ★                    | GNSSR 2022-05-22 15:27:29     |
| ■                    | ICESat-2 23-Feb-2022 21:56:25 |
| ■                    | ICESat-2 23-Nov-2022 08:55:57 |
| ■                    | ICESat-2 22-Aug-2023 19:54:32 |
| ■                    | ICESat-2 21-Nov-2023 15:34:24 |

Importance: Without removing stable structures, comparing water height measurements collected at different regions of a lake may lead to biased results

# Summary

1. **Good agreement across ICESat-2, GNSS-R, and SWOT**


- Started with case-studies comparing data collected in similar place & time

2. **Extended analysis over the same location to a diverse range of times to understand the result, which strongly suggests a geoid anomaly**

- 9 SWOT
- 2 GNSS-R
- 4 ICESat-2

3. **Moving forward:**

- Expand to studying more of the lake
- Determine best way to combine measurements and separate seasonal signatures from geoid signatures

Mission	Advantages	Challenges
ICESat-2	<ul style="list-style-type: none"> <li>• Lowest noise over open water</li> <li>• Dense photon cloud w/in a track</li> <li>• 3-6 laser tracks in 1 pass</li> </ul>	<ul style="list-style-type: none"> <li>• Fewer total number of useable tracks (clouds)</li> <li>• Strongly influenced by vegetation</li> </ul>
GNSS-R	<ul style="list-style-type: none"> <li>• Low-cost</li> <li>• Many tracks, diverse orientations</li> <li>• Penetrates vegetation, clouds</li> </ul>	<ul style="list-style-type: none"> <li>• Relative nature of carrier-phase altimetry</li> </ul> 
SWOT	<ul style="list-style-type: none"> <li>• Swath data</li> <li>• Frequent repeat</li> </ul>	<ul style="list-style-type: none"> <li>• Influenced by emergent riparian vegetation</li> <li>• Ka-band attenuated by rain</li> </ul>

# Acknowledgements

- **This work funded by NASA #80NSSC22K1116, #80NSSC21K1011, and NASA FINESST #80NSSC23K1554**
- **Spire data made accessible through the NASA Commercial SmallSat Data Acquisition (CSDA) program**
- **A portion of this work was carried out at the Jet Propulsion Laboratory, California Institute of Technology under a contract with the National Aeronautics and Space Administration**

# Data References

- **ICESat-2 ATL03:** T. A. Neumann, A. Brenner, D. Hancock, J. Robbins, J. Saba, K. Harbeck, A. Gibbons, J. Lee, S. B. Luthcke, T. Rebold, et al, “ATLAS/ICESat-2 L2A Global Geolocated Photon Data, Version 5 [Data Set]”. *Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center*, 2021. <https://doi.org/10.5067/ATLAS/ATL03.005>
- **SWOT:** JPL D-105504, SWOT Algorithm Theoretical Basis Document: Level 2 KaRIn High Rate Pixel Cloud Science Algorithm Software, *Jet Propulsion Laboratory Internal Document*. (2023)

JPL D-105501, SWOT Algorithm Theoretical Basis Document: Level 2 KaRIn High Rate Raster (L2 HR Raster) Science Algorithm Software, *Jet Propulsion Laboratory Internal Document*. (2023).

**Water Mask:** J.F. Pekel, A. Cottam, N. Gorelick, and A. S. Belward, “High-resolution mapping of global surface water and its long term changes,” *Nature*, vol. 540, pp. 418- 422, 2016

• **Data access:** *EC JRC/Google*

- **DAHITI:** C. Schwatke, D. Dettmering, W. Bosch, and F. Seitz, “Dahiti– an innovative approach for estimating water level time series over inland waters using multi-mission satellite altimetry,” *Hydrol. Earth Syst. Sci.*, vol. 19, pp. 4345–4364, 2015.






**JPL**

# Back-up Slides



# Water Mask from Global Water Surface Explorer

Pekel, 2016 (doi:10.1038/nature20584). Source: EC JRC/Google

Landsat-derived Occurrence from 1984 - 2021




# GNSS-R Track Comparisons



# Data Match-up Strategy

- **Compare tracks within:**
  - 3 km separation (max)
  - 2 weeks



University of Colorado



Motivation






Methods

Results

Summary

# Definition of Circular Length

$$\zeta = \frac{1}{N} \left| \sum_1^N \cos \delta\phi_i + j \sum_1^N \sin \delta\phi_i \right|, \quad \zeta \in [0, 1]$$



Higher  $\zeta$  indicates a more coherent phase